

V. V. Buyadzhi

Odessa State Environmental University, L'vovskaya str.15, Odessa-16, 65016, Ukraine
E-mail: vbuyad@mail.ru

LASER MULTIPHOTON SPECTROSCOPY OF ATOM EMBEDDED IN DEBYE PLASMAS: MULTIPHOTON RESONANCES AND TRANSITIONS

The consistent relativistic energy approach to atom in a realistic laser field, based on the Gell-Mann and Low S-matrix formalism, is applied to studying the resonant multiphoton transitions in atoms embedded in the Debye plasmas. There is considered a new scheme to calculating the multiphoton transitions characteristics, shifts and widths of multiphoton resonances. An approach is used for treating the three-photon transitions in krypton embedded in the Debye plasmas.

1. The physics of multiphoton phenomena is one of the very quickly developed branches of the modern optics and spectroscopy, photophysics. Studying of multiphoton phenomena in atoms, molecules etc has a great progress that is stimulated by development of new laser technologies (see Refs. [1-10]). The appearance of the powerful laser sources allowing to obtain the radiation field amplitude of the order of atomic field in the wide range of wavelengths results to systematic investigations of the nonlinear interaction of radiation with atomic and molecular systems [1-14].

At the same time a direct laser-nucleus interactions traditionally have been dismissed because of the well known effect of small interaction matrix elements [9-11]. Some exceptions such as an interaction of x-ray laser fields with nuclei in relation to alpha, beta-decay and x-ray-driven gamma emission of nuclei have been earlier considered. With the advent of new coherent x-ray laser sources in the near future, however, these conclusions have to be reconsidered.

At present time a great interest has been connected with studying atomic processes in plasma environments because of the plasma environment screening effect on the plasma-embedded atomic systems. One should remind that the screening effects have play a important and significant part in the investigation of plasma environments over the past several decades.

Different theoretical methods have been employed along with the Debye screening to study plasma environments.

The interaction of atoms with the external alternating fields, in particular, laser fields, has been the subject of intensive experimental and theoretical studied (see, for example, Refs. [1-8, 12-24]). A definition of the k-photon emission and absorption probabilities and atomic levels shifts, study of dynamical stabilization and field ionization etc are the most actual problems to be solved.

Above methods which are usually used one should mention such approaches as the standard perturbation theory (surely for low laser field intensities), Green function method, the density-matrix formalism, time-dependent density functional formalism, direct numerical solution of the Schrödinger (Dirac) equation, multi-body multiphoton approach, the time-independent Floquet formalism etc (see [1-8,12-24] and Refs. therein).

Earlier the relativistic energy approach to studying the interaction of atom with a realistic strong laser field, based on the Gell-Mann and Low S-matrix formalism, has been developed. Originally, Ivanov has proposed an idea to describe quantitatively a behaviour of an atom in a realistic laser field by means studying the radiation emission and absorption lines and further the theory of interaction of an atom with the Lorenz laser pulse and calculating the corresponding

lines moments has been in details developed in Ref. [19-25]. It has been checked in numerical simulation of the multiphoton resonances shifts and widths in the hydrogen and caesium. Theory of interaction of an atom with the Gauss and soliton-like laser pulses and calculating the corresponding lines moments has been in details presented in Refs. [23,26,27].

Here the consistent relativistic energy approach to atom in a realistic laser field, based on the Gell-Mann and Low S-matrix formalism, is applied to studying the resonant multiphoton transitions in atoms embedded in the Debye plasmas. There is considered a new scheme to calculating the multiphoton transitions characteristics, shifts and widths of multiphoton resonances. An approach is used for treating the three-photon transitions in krypton embedded in the Debye plasmas

2. The relativistic energy approach in the different realizations and the radiation lines moments technique is in details presented in Refs. [19-30]. So, here we are limited only by presenting the master elements. In the theory of the non-relativistic atom a convenient field procedure is known for calculating the energy shifts dE of degenerate states. This procedure is connected with the secular matrix M diagonalization. In constructing M , the Gell-Mann and Low adiabatic formula for dE is used [20-23,31]. In relativistic theory, the Gell-Mann and Low formula dE is connected with electro-dynamical scattering matrix, which includes interaction with as a laser field as a photon vacuum field. A case of interaction with photon vacuum is corresponding to standard theory of radiative decay of excited atomic states. Surely, in relativistic theory the secular matrix elements are already complex in the second perturbation theory (PT) order. Their imaginary parts are connected with radiation decay possibility. The total energy shift is usually presented in the form [23]:

$$\delta E = \text{Re}\delta E + i \text{Im}\delta E ,$$

$$\text{Im} \delta E = -P/2, \quad (1)$$

where P is the level width (decay possibility). Let us describe the interaction “atom-laser field” by the Ivanov potential [21,23]:

$$V(r,t)=V(r)\int d\omega f(\omega-\omega_0) \sum_{n=-\infty}^{\infty} \cos[\omega_0 t + \omega_0 n \tau] \quad (2)$$

Here ω_0 is the central laser radiation frequency, n is the whole number. The function $f(\omega)$ is a Fourier component of the laser pulse. The condition $\int d\omega f^2(\omega)=1$ normalizes potential $V(r,t)$ on the definite energy in the pulse. Usually one could consider the pulses with Lorentz shape (coherent 1-mode pulse): $f(\omega) = b/(\omega^2+D^2)$, Gaussian one (multi-mode chaotic pulse): $f(\omega) = b \exp[\ln 2(\omega^2/D^2)]$ and the soliton-like pulse: $f(t) = b \text{ch}^{-1}[t/D]$ (b -normalizing multiplier).

The master program results in the calculating an imaginary part of energy shift $\text{Im}dE_a(\omega_0)$ for any atomic level as the function of the central laser frequency ω_0 . An according function has the shape of the resonance, which is connected with the transition $a-p$ (a , p -discrete levels) with absorption (or emission) of the “ k ” number of photons. For this transition the following values are determined [20-23]:

$$\delta\omega(p\alpha|k)=\int' d\omega \text{Im} \delta E_\alpha(\omega)(\omega - \omega_{p\alpha}/k)/N, \quad ($$

$$\mu_m = \int' d\omega \text{Im} \delta E_\alpha(\omega) (\omega - \omega_{p\alpha}/k)^m / N,$$

where

$$\int' d\omega \text{Im} E_\alpha$$

is the normalizing multiplier; ω_{pa} is position of the non-shifted line for transition $a-p$, $d\omega(p\alpha|k)$ is the line shift under k -photon absorption; $\omega_{pa} = \omega_{pa} + k \times d\omega(p\alpha|k)$. The first moments m_1 , m_2 and m_3 determine the atomic line centre shift, its dispersion and the asymmetry.

To find m_m , we need to get an expansion of E_a to PT series:

$$E_\alpha = \sum E_\alpha^{(2k)}(\omega_0).$$

One may use here the Gell-Mann and Low adiabatic formula for dE_a [20-23]. The consideration can be simplified by account of the k -photon absorption contribution in the first two PT orders. Besides, summation on laser pulse is exchanged by integration. The corresponding $(l+2k+1)$ -times integral on $(l+2k)$ temporal variables and r ($l=0,2$) (integral I_g) are calculated [19-23]. Finally, after some cumbersome transformations one can get the expressions for the line moments. The corresponding expressions for the Gaussian laser pulse are as follows:

$$\delta\omega(p\alpha | k) = \{\pi\Delta/(k+1)k\} [E(p, \omega_{p\alpha}/k) - E(\alpha, \omega_{p\alpha}/k)], \quad (9)$$

$$\mu_2 = \Delta^2/k$$

$$\mu_3 = \{4\pi\Delta^3/[k(k+1)]\} [E(p, \omega_{p\alpha}/k) - E(\alpha, \omega_{p\alpha}/k)],$$

where

$$E(j, \omega_{p\alpha}/k) = 0,5 \sum_{p_i} V_{jpi} V_{pij} \left[\frac{1}{\omega_{jpi} + \omega_{p\alpha}/k} + \frac{1}{\omega_{jpi} - \omega_{p\alpha}/k} \right] \quad (10)$$

The summation in (10) is over all atomic states. Let us note that these formulas for the Gaussian pulse differ of the Lorenz shape laser pulse expressions [21-23]. For the soliton-like pulse it is necessary to carry out the numerical calculation or use some approximations to simplify the expressions [27].

In order to calculate (10), one should use the technique [28,29] of calculating sums of the QED PT second order, which has been earlier applied by us in calculations of some atomic and mesoatomic parameters [26,27,30-32].

Finally the computational procedure results in a solution of the ordinary differential equations system for above described functions and integrals. In concrete numerical calculations the PC "Superatom-ISAN" package is used. The construction of the operator wave functions bases within the QED PT, the technique of calculating the matrix elements in Eqs. (9,10) and other details is presented in Refs. [19-30].

3. In order to take into account the plasmas screening effect one could use the known Debye shielding model. As it is well known (c.f.[33-35] and refs there) in the classical theory of plasmas developed by Debye and Hückel, the interaction potential between two charged particles in a plasma is modelled by a Yukawa-type potential as follows:

$$V(r_i, r_j) = (Z_a Z_b / |r_a - r_b|) \exp(-\mu |r_a - r_b|), \quad (11)$$

where r_a, r_b represent respectively the spatial coordinates of particles A and B and Z_a, Z_b denote their charges. A difference between the Yukawa

type potential and standard Coulomb potential is in account for the effect of plasma, which is modeled by the shielding parameter m [33]. The parameter m is linked with the plasma parameters such as the temperature T and the charge density n as follows $\mu \sim \sqrt{e^2 n / k_B T}$ where, as usually, e is the electron charge and k_B is the Boltzman constant. The density n is given as a sum of the electron density N_e and the ion density N_k of the k -th ion species having the nuclear charge q_k . Let us remind [35] that under typical laser plasma conditions of $T \sim 1$ keV and $n \sim 10^{23} \text{ cm}^{-3}$ the parameter m is of the order of 0,1 in atomic units. By introducing the Yukawa-type electron-nuclear attraction and electron-electron repulsion potentials, the electronic Hamiltonian for N -electron multicharged ion in a plasma is given in atomic units as follows:

$$H = \sum_i [\alpha p_i^2 - \beta m_i^2 - Z \exp(-\mu r_i) / r_i] + \sum_{i>j} \frac{(1 - \alpha_i \alpha_j)}{r_j} \exp(-\mu r_j) \quad (12)$$

A difference between the Hamiltonian (12) and analogous model Hamiltonian with the Yukawa potential of ref. [33] is in using the relativistic approximation, which is obviously necessary for adequate description of relativistic systems [35].

4. In ref. [36] there were presented the results of the numerical simulation for the three-photon resonant, four-photon ionization profile of atomic krypton (the $4p \text{ @ } 5d[1/2]_1$ and $4p \text{ @ } 4d[3/2]_1$ three photon Kr resonances are considered; intense uv (285-310 nm) laser radiation with intensity range $3 \cdot 10^{12} - 10^{14} \text{ W/cm}^2$ studied) in a free state (n in sense of the absence a plasmas environment). There have been determined the corresponding parameters of the $4p \text{ @ } 5d[1/2]_1$ (i) and $4p \text{ @ } 4d[3/2]_1$ (ii) three photon Kr resonances. The resonance shift is proportional to intensity with a width dominated by lifetime broadening of the excited state.

The numerical simulation [36] results for the $4p \text{ @ } 5d[1/2]_1$ (i) and $4p \text{ @ } 4d[3/2]_1$ (ii) three photon Kr resonances are as follows: (i) the shift $dw_0(pa|3) = aI$, $a_{\text{exp}} = 3.95 \text{ meV}/(\text{Tw} \times \text{cm}^{-2})$ and width $b_{\text{exp}} = 1.5 \text{ meV}/(\text{Tw} \times \text{cm}^{-2})$; (ii) shift $dw_0(pa|3) = aI$, $a_{\text{exp}} = 8.1 \text{ meV}/(\text{Tw} \times \text{cm}^{-2})$ and width $b_{\text{exp}} = 4.2 \text{ meV}/(\text{Tw} \times \text{cm}^{-2})$. We have chosen the Debye length parameter values $\lambda_D = 50$ (25) and the corresponding computed coefficients are as

follows: (i) $a=3.76 \text{ meV}/(\text{Tw}\times\text{cm}^{-2})$ ($a=3.2 \text{ meV}/(\text{Tw}\times\text{cm}^{-2})$); (ii) $a=7.8 \text{ meV}/(\text{Tw}\times\text{cm}^{-2})$ ($a=6.5 \text{ meV}/(\text{Tw}\times\text{cm}^{-2})$).

The presented results show that Debye plasma environments have an effect on the multiphoton transitions. Nevertheless, one should keep in mind some important facts the (see, for example, [33,34]). It is clear the static screening result considered above is subject to the condition that the plasma is a thermodynamically equilibrium plasma and neglects the contributions from ions in plasma since electrons provide more effective shielding than ions.

Obviously with the changing the plasma conditions (parameters) in principle there can be taken a place a significant variations. Besides, one should remember about the conditions of applicability of the Debye approximation.

References

1. Batani D., Joachain C. J., eds. 2006 Matter in super-intense laser fields, AIP Serie, N.-Y.
2. Ivanova E. P. 2011 Phys.Rev.A 84 043829
3. Jaron-Becker A., Becker A., Faisal F. H. M 2004 Phys.Rev.A 69 023410
4. Plummer M, Noble C J 2003 J.Phys. B: At. Mol. Opt. Phys. 36 L219
5. van der Hart H. W., Lysaght M. A., Burke P. G. 2007 Phys.Rev.A 76 043405
6. Mercouris T., Nikolaidis C. A. 2003 Phys. Rev. A 67 063403
7. Brandas E., Floelich P 1977 Phys. Rev. A 16 2207
8. Kwanghsi Wangt, Tak-San Hot, Shih-I Chu 1985 J.Phys. B: At. Mol. Opt. Phys. 18 4539; 2006 Phys.Rev.A 73 023403
9. Glushkov A. V., Ivanov L. N., Letokhov V. S. Nuclear quantum optics 1991 Preprint of Institute for Spectroscopy, USSR Academy of Sciences, Troitsk-N7
10. Bürvenich T. J., Evers J., Keitel C. H. 2006 Phys.Rev.Lett. 96 142501; 2006 Phys.Rev.C 74 044601
11. Shahbaz A., Müller C., Staudt A., Bürvenich T. J., Keitel C. H. 2007 Phys. Rev.Lett. 98 263901
12. Fedorov M. V. 1999 Physics-Uspekhi 169 66
13. Delone N. B., Kraynov V. P. 1998 Physics-Uspekhi 168 531
14. Belyaev V. S., Kraynov V. P., Lisitsa V. S., Matafonov A. P. Physics-Uspekhi 178 823
15. Zoller P. 1982 J.Phys. B: At. Mol. Opt. Phys. 15 2911
16. Lompre L-A., Mainfrau G., Manus C., Marinier J. P. 1981 J.Phys. B: At. Mol. Opt. Phys. 14 4307
17. Khetselius O.Yu., 2012 J. of Phys.: Conf. Ser.. 397 012012; 2009 Phys.Scr. T.135 014023.
18. Landen O. L., Perry M. D., Campbell E. M. 1987 Phys.Rev.Lett. 59 2558
19. Ivanov L. N., Letokhov V. S. 1975 JETP 68 1748; 1985 Com. Mod. Phys. D4 169
20. Ivanov L. N., Ivanova E. P., Aglitsky E. V. 1988 Phys. Rep. 166 315
21. Glushkov A. V., Ivanov L. N., 1985 Preprint of ISAN, USSR Academy of Sci., Troitsk, N3.
22. Glushkov A. V., Ivanov L. N., Ivanova E. P., 1986 Autoionization Phenomena in Atoms, Moscow Univ. Press, Moscow, p 58
23. Glushkov A. V., Ivanov L. N. 1992 Phys. Lett. A 170 33.
24. Glushkov A. V., Ivanov L. N. 1992 Proc. of 3rd Seminar on Atomic Spectroscopy (Chernogolovka, Moscow reg.) p 113
25. Glushkov A, Ivanov L N 1993 J.Phys. B: At. Mol. Opt. Phys. 26 L179.
26. Glushkov A. V., Loboda A., Gurnitskaya E., Svinarenko A., 2009 Phys. Scripta T135 014022
27. Glushkov A. V., Khetselius O. Yu., Loboda A. V., Svinarenko A. A., 2008 Frontiers in Quantum Systems in Chem. and Phys., Ser. Progress in Theoretical Chemistry and Physics, vol. 18. ed. S Wilson, P. J., Grout, J., Maruani, G.; Delgado-Barrio, P. Piecuch (Berlin: Springer) p 543
28. Ivanov L. N., Ivanova E. P., 1974 Theor. Math.Phys. 20 282
29. Ivanova E P, Ivanov L N, Glushkov A V, Kramida A E 1985 Phys. Scripta 32 512

30. Glushkov A V 2012 Advances in the Theory of Quantum Systems in Chemistry and Physics. Ser. Progress in Theoretical Chemistry and Physics, vol. 28, ed. by K Nishikawa, J Maruani, E Brandas, G Delgado-Barrio, P Piecuch (Berlin: Springer) p 131
31. Glushkov A V, Khetselius O Yu, Malinovskaya S V 2008 Europ. Phys. Journ ST 160 195.
32. Khetselius O Yu, 2009 Int. Journ. Quant. Chem. 109 3330 ;
33. Okutsu, H.; Sako, T.; Yamanouchi, K.; Dierksen, G.H.F. J. Phys. B: At. Mol. Opt. Phys. 2005, 38, 917-928; Badnell, N.R. Journ. Phys. CS. 2007, 88, 012070 (8p.).
34. Paul S., Ho Y.K., J. Phys. B: At. Mol. Opt. Phys. 2010, 43, 065701
35. Glushkov, A.V.; Malinovskaya, S.V.; Prepelitsa, G.P.; Ignatenko, A.V. et al Journ. Phys. CS. 2004, 11, 199–208; Int.J. Quant. Chem. 2011, 111,288-298.
36. Nakamura, N. ; Kavanagh, A.P. ; Watanabe, H. ; Sakaue, H.A. ; Li, Y.; Kato, D. ; Curell, F.J. ; Ohtani S. Journ. Phys. CS. 2007, 88, 012066 (6p.); Li, Y.; Wu, J. ; Hou, Y.; Yuan, J. J. Phys. B: At. Mol. Opt. Phys. 2008, 41, 145002 (8p.).
37. Buyadzhi V.V., Glushkov A.V., Lovett L., Photoelectronics. 2014, 23, 38-43.

This article has been received within 2015

UDC 539.182

V. V. Buyadzhi

LASER MULTIPHOTON SPECTROSCOPY OF ATOM EMBEDDED IN DEBYE PLASMAS: MULTIPHOTON RESONANCES AND TRANSITIONS

Abstract.

The consistent relativistic energy approach to atom in a realistic laser field, based on the Gell-Mann and Low S-matrix formalism, is applied to studying the resonant multiphoton transitions in atoms embedded in the Debye plasmas. There is considered a new scheme to calculating the multiphoton transitions characteristics, shifts and widths of multiphoton resonances. An approach is used for treating the three-photon transitions in krypton embedded in the Debye plasmas.

Key words: electromagnetic interactions, laser field, multiphoton resonances, plasmas

УДК 539.182

В. В. Буюджи

ЛАЗЕРНАЯ МУЛЬФОТОННАЯ СПЕКТРОСКОПИЯ АТОМОВ В ДЕБАЕВСКОЙ ПЛАЗМЕ : МНОГОФОТОННЫЕ ПЕРЕХОДЫ И РЕЗОНАНСЫ

Резюме.

Релятивистский энергетический подход к описанию спектроскопии атома в лазерном поле, основывающийся на S-матричном формализме Гелл-Манна и Лоу, применяется для изучения

резонансных многофотонных переходов атомов в дебаевской плазме. Предложена новая схема вычисления характеристик многофотонных переходов, энергий и ширин многофотонных переходов. Подход использован для описания трехфотонных переходов в криптоне в плазме Дебая.

Ключевые слова: электромагнитное, лазерное поле, многофотонные резонансы, плазма

УДК 539.182

В. В. Буяджи

ЛАЗЕРНА МУЛЬФОТОНА СПЕКТРОСКОПІЯ АТОМІВ У ДЕБАЄВСЬКІЙ ПЛАЗМІ: БАГАТОФОТОННІ ПЕРЕХОДИ І РЕЗОНАНСИ

Резюме.

Релятивістський енергетичний підхід до опису спектроскопії атома в лазерному полі, що ґрунтується на S-матричному формалізмі Гелл-Манна і Лоу, застосовується для вивчення резонансних багато фотонних переходів атомів в дебаєвській плазмі. Запропоновано нову схему обчислення характеристик багатофотонних переходів, енергій і ширин багато фотонних переходів. Підхід використаний для опису трьох фотонних переходів у криптоні в плазмі Дебая.

Ключові слова: електромагнітна взаємодія, лазерне поле, багатофотонні резонанси, плазма